

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/350311889>

Effects of Modified Biochars on the Growth of Maize (*Zea mays* L.)

Poster · September 2016

DOI: 10.13140/RG.2.2.33944.52484

CITATIONS

0

READS

937

8 authors, including:



Md Arifur Rahaman

Chinese Academy of Agricultural Sciences

22 PUBLICATIONS 169 CITATIONS

SEE PROFILE



Sajid Latif

75 PUBLICATIONS 4,257 CITATIONS

SEE PROFILE



Kiatkamjon Intani

Kasetsart University

16 PUBLICATIONS 259 CITATIONS

SEE PROFILE



Joachim Müller

University of Hohenheim

535 PUBLICATIONS 9,939 CITATIONS

SEE PROFILE

Charlotte C. Dietrich¹, Arifur Rahaman^{1,2}, Sajid Latif², Kiatkamjon Intani², Joachim Müller² and Nicolai D. Jablonowski¹

¹ Forschungszentrum Jülich, IBG-2 Plant Sciences; ² Universität Hohenheim, Institute of Agricultural Engineering, Tropics and Subtropics Group
c.dietrich@fz-juelich.de; s.latif@uni-hohenheim.de

Objectives

- Evaluate the fertilizing effect of (modified) biochars since biochar applications have been shown to either elevate plant productivity or impair soils' nutrient status.
- Identify biochar induced effects on plant functional traits and plant productivity to derive mechanistic knowledge required to develop tailor-made chars.
- Observe the effect of a possible nitrogen immobilization following char application, as certain biochar compounds are hypothesized to reduce nitrogen availability.
- We hypothesize that biochar application affects i) soil pH and nutrient availabilities and thus ii) plant productivity and traits.


Method

- Biochar was produced from whole maize cobs at University of Hohenheim using a self-purging pyrolysis reactor. The biochar was modified using i) washing (with either ethanol or HCl) and ii) loading (with digestate, a nutrient-rich byproduct of the biogas production) processes. Non-modified char served as the positive control.
- Maize was grown on a sand/biochar mixture for a period of 35 days. Unamended nutrient-deficient substrate served as the negative control.




Results


Shoot biomass: highest yield observed with loaded biochar



Plant productivity: loaded biochar doubled total biomass production



C/N ratio (plant tissue): loaded biochar decreased C/N ratio



Relative biochar induced changes on above ground biomass (left), total biomass production (middle) and plant C/N ratio (right) in dependence of the unamended control. We performed separate non-parametric ANCOVA's using Euclidian distance measures with time as a covariate to assess how biochar amendment affects plant functional parameters. We took 8 replicates at three harvests each (21, 28 and 35 days of plant growth). Small letters indicate significant differences ($p < 0.05$) between groups (small letter "a" indicates differences between treatment and the unamended control). The η^2 values indicate the relative effect sizes of each explaining factor on plant parameter variation. Data is shown as means and their standard error.

- Biochar applications increased total C and N concentrations in all mixtures. Plant-availability of Mg was significantly reduced in non-modified biochar, ethanol-washed biochar and loaded biochar.
- Biochar amendments affected plant functional traits, as all chars induced a significant increase in above ground biomass production.
- Nutrient loaded biochar enhanced nitrogen uptake while other biochars reduced or maintained nitrogen uptake relative to those of the control treatments. In consequence, C/N ratios in plant tissue tended to increase for all biochar treatments except digestate loaded chars, which induced a significant decline of C/N ratios.
- Nutrient loaded biochar doubled biomass production of emerged maize plants, although soil N content did not significantly increase among treatments.

Discussion

- Our results suggest that plant productivity remained unchanged after application of biochar and its washed forms. Contrastingly, nutrient loaded biochar induced a significant increase in productivity due to markedly improved plant nutrient uptake that affected biomass productivity and allocation.
- This experiment indicates that establishing loaded chars as efficient nutrient carriers might circumvent N-immobilization processes associated with an impaired C/N balance following char application.
- Future research is required to develop biochar concepts that facilitate prolonged improvements of plant productivity of emerged maize plants.